Cumulative Author Index, Volume 17

A

Abramson DL, Pribaz JJ, Orgill DP. The use of free tissue transfer in burn reconstruction. 1996;17:402-8
Akman N (*see* Haberal et al). 1996;17:147-9
Archer SB (*see* Brown et al). 1996;17:552-7
Atiles L (*see* Spann et al). 1996;17:514-7
Auger FA (*see* Xu et al). 1996;17:7-13

B

Baker RAU, Jones S, Sanders C, Sadinski C, Martin-Duffy K, Berchin H, Valentine S. Degree of burn, location of burn, and length of hospital stay as predictors of psychosocial status and physical functioning. 1996;17:327-33
Bala RJ (*see* McMillen et al). 1996;17:384-9
Barese S (*see* Klein et al). 1996;17:565-70
Barillo DJ, Goode R. Substance abuse in victims of fire. 1996;17:71-6
—— (*see* Harvey et al). 1996;17:176-80
Barrash J (*see* Janus and Barrash). 1996;17:409-15
Başgöze O (*see* Haberal et al). 1996;17:147-9
Benaim F (*see* Patiño et al). 1996;17:528-31
Ben-Bassat H, Strauss N, Ron M, Chaouat M, Breiterman S, Israeli A, Wexler MR, Eldad A. Transplantation performance of human skin cryopreserved by programmed or stepwise freezing and stored at -80°C or -180°C. 1996;17:421-8
Bentrem DJ (*see* Kirby et al). 1996;17:156-60
Berchin H (*see* Baker et al). 1996;17:327-33
Berger N (*see* Turner et al). 1996;17:169-74
Bettinger DA, Mast B, Gore D. Hyaluronic acid impedes reepithelialization of skin graft donor sites. 1996;17:302-4
Bilevich ED (*see* Davis et al). 1996;17:558-61
Biondo N (*see* Klein et al). 1996;17:518-21
Blakeney P (*see* LeDoux et al). 1996;17:472-6
—— (*see* Moore et al). 1996;17:547-51
Blank V (*see* Jenkins et al). 1996;17:49-61
Bolgiani A (*see* Patiño et al). 1996;17:528-31
Bradrick JP (*see* Punjabi et al). 1996;17:285 (Letter)
Breiterman S (*see* Ben-Bassat et al). 1996;17:421-8
Brigham PA, McLoughlin E. Burn incidence and medical care use in the United States: estimates, trends, and data sources. 1996;17:95-107
Britton SM. Proper use of nonadherent impregnated dressings. 1996;17:93-4 (Letter)
Brown DL, Archer SB, Greenhalgh DG, Washam MA, James LE, Warden GD. Inhalation injury severity scoring system: a quantitative method. 1996;17:552-17
Brown RL, Henke A, Greenhalgh DG, Warden GD. The use of haloperidol in the agitated, critically ill pediatric patient with burns. 1996;17:34-8
Burger M (*see* Kealey and Burger). 1996;17:S6-7

C

Cadier MA, Clarke JA. Dermasorb versus Jelonet in patients with burns skin graft donor sites. 1996;17:246-51
Caldwell FT Jr, Wallace BH, Cone JB. Sequential excision and grafting of the burn injuries of 1507 patients treated between 1967 and 1986: end results and the determinants of death. 1996;17:137-46
Calvano SE (*see* Lawrence et al). 1996;17:1-6
Caruso DM, Gregory MW, Schiller WR. The use of skin from a monozygotic twin combined with cultured epithelial autografts as coverage for a large surface area burn: a case report and review of the literature. 1996;17:432-4 (Case rep.)
Cassuto J (*see* Tarnow et al). 1996;17:207-12
Cazzaniga AL (*see* Davis et al). 1996;17:558-61
Chaouat M (*see* Ben-Bassat et al). 1996;17:421-8
Chapo GA (*see* Klein et al). 1996;17:565-70
Cioffi WG (*see* Harvey et al). 1996;17:176-80
Clarke JA (*see* Cadier and Clarke). 1996;17:246-51
Cone JB (*see* Caldwell et al). 1996;17:137-46
Connolly ML (*see* Dattolo et al). 1996;17:182-7
Cook EL (*see* Watkins et al). 1996;17:78-92
Cooper ML, Hansbrough JF, Polarek JW. The effect of an arginine–glycine–aspartic acid peptide and hyaluronate synthetic matrix on epithelialization of meshed skin graft interstices. 1996;17:108-16
Cornwell P. Critical pathways in the QI process. 1996;17:S3-6
Covington DS, Wainwright DJ, Parks DH. Prognostic indicators in the elderly patient with burns. 1996;17:222-30
Cunningham ME (*see* McMillen et al). 1996;17:384-9
Cuono CB (*see* Klein et al). 1996;17:565-70

D

D'Aiuto ML (*see* McMillen et al). 1996;17:384-9
Dattolo J, Trout S, Connolly ML. Home health care and burn care: an educational and economical program. 1996;17:182-7
Davis SC, Mertz PM, Bilevich ED, Cazzaniga AL, Eaglstein WH. Early debridement of second-degree burn wounds enhances the rate of epithelization—an animal

*January/February, pp. 1-94; March/April, pp. 95-190; May/June, pp. 191-286; July/August, pp. 287-378; September/October, pp. 379-476; November/December, Part 1, 477-570. November/December, Part 2, pp. S1-S36.

model to evaluate burn wound therapies. 1996;17:558-61
—— (*see* Mertz et al). 1996;17:199-206
DeBiasse MA (*see* Gamliel et al). 1996;17:264-72
DeClement FA (*see* Hughes et al). 1996;17:243-5
Del Rio JL (*see* Holguin et al). 1996;17:150-4
Demling R (*see* LaLonde et al). 1996;17:379-83
Demling RH (*see* Gamliel et al). 1996;17:264-72
de Riesthal HF (*see* Lawrence et al). 1996;17:1-6
Desai MH (*see* Mukhdomi et al). 1996;17:346-50
deSerres S (*see* Tabata et al). 1996;17:231-6
Dimick A (*see* Wainwright et al). 1996;17:124-36
Dissanayake S (*see* Doong et al). 1996;17:497-514
Doong H, Dissanayake S, Gowrishankar TR, LaBarbera MC, Lee RC. Calcium antagonists alter cell shape and induce procollagenase synthesis in keloid and normal human dermal fibroblasts. 1996;17:497-514
Dries DJ (*see* Stengle et al). 1996;17:14-8
Duncan JW (*see* Still et al). 1996;17:562-4
Dunnick CA, Gibran NS, Heimbach DM. Substance P has a role in neurogenic mediation of human burn wound healing. 1996;17:390-6
Dutcher KD (*see* Fricke et al). 1996;17:338-45

E

Eaglstein WH (*see* Davis et al). 1996;17:568-61
Edlich RF. Clarification on the use of silver-based purification systems. 1996;17:285 (Letter)
—— (*see* Kirby et al). 1996;17:156-60
Eldad A (*see* Ben-Bassat et al). 1996;17:421-8
Engrav LH (*see* Fricke et al). 1996;17:338-45

F

Fitzpatrick JC (*see* Harvey et al). 1996;17:176-80
Foertsch CE, O'Hara MW, Stoddard FJ, Kealey GP. Parent participation during burn debridement in relation to behavioral distress. 1996;17:372-7
Franzen L (*see* Mertz et al). 1996;17:199-206
Fratianne RB (*see* Yurko and Fratianne). 1996;17:S2-3
Fraulin FOG, Illmayer SJ, Tredget EE. Assessment of cosmetic and functional results of conservative versus surgical management of facial burns. 1996;17:19-29
Fricke NB, Omnell L, Dutcher KD, Hollender LG, Engrav LH. Skeletal and dental disturbances after facial burns and pressure garments. 1996;17:338-45
Frye KE, Izenberg SD, Williams MD, Luterman A. Simulated biologic intelligence used to predict length of stay and survival of burns. 1996;17:540-6

G

Gamliel Z, DeBiasse MA, Demling RH. Essential microminerals and their response to burn injury. 1996;17:264-72
Garcia JP (*see* Holguin et al). 1996;17:150-4
Garofalo RP (*see* Zimmer et al). 1996;17:491-6
Germain L (*see* Xu et al). 1996;17:7-13
Gibran NS (*see* Dunnick et al). 1996;17:390-6
Goode R (*see* Barillo and Goode). 1996;17:71-6
Gordon M. Introduction. 1996;17:S1
Gordon M. Preface. 1996;17:S2
Gore D (*see* Bettinger et al). 1996;17:302-4
—— (*see* Sanford and Gore). 1996;17:323-6
Goulet F (*see* Xu et al). 1996;17:7-13
Gowrishankar TR (*see* Doong et al). 1996;17:497-514
Grana D (*see* Patiño et al). 1996;17:528-31
Greenhalgh DG (*see* Brown et al). 1996;17:34-8
—— (*see* Brown et al). 1996;17:552-7
Gregory MW (*see* Caruso et al). 1996;17:432-4 (Case rep.)
Greulich KM (*see* Young et al). 1996;17:305-10
Griswold JA (*see* Penturf et al). 1996;17:465-70
Grobmyer SR, Maniscalco SP, Purdue GF, Hunt J. Alcohol, drug intoxication, or both at the time of burn injury as a predictor of complications and mortality in hospitalized patients with burns. 1996;17:532-9
Gudjonsson O (*see* Yelon et al). 1996;17:241-2 (Case rep.)
Gupta A (*see* Sjöberg and Gupta). 1996;17:285-6 (Letter)
Gürer S (*see* Haberal et al). 1996;17:147-9

H

Haberal MA, Gürer S, Akman N, Başgöze O. Persistent peripheral nerve pathologies in patients with electric burns. 1996;17:147-9
Hansbrough JF (*see* Cooper et al). 1996;17:108-16
—— (*see* Rennekampff et al). 1996;17:213-21
—— (*see* Rennekampff et al). 1996;17:522-7
Harries T (*see* Linder et al). 1996;17:351-2 (Case rep.)
Harvey KD, Barillo DJ, Hobbs CL, Mozingo DW, Fitzpatrick JC, Cioffi WG, McManus WF, Pruitt BA Jr. Computer-assisted evaluation of hand and arm function after thermal injury. 1996;17:176-80
Hastings B, Roth A, Nolan D, Miller S. Wound coverage: Is there a difference? 1996;17:416-20
Heimbach D (*see* Wainwright et al). 1996;17:124-36
Heimbach DM (*see* Dunnick et al). 1996;17:390-6
Henke A (*see* Brown et al). 1996;17:34-8
Hennigan J (*see* LaLonde et al). 1996;17:379-83
Hensell DO (*see* Hughes et al). 1996;17:243-5
Herndon D (*see* LeDoux et al). 1996;17:472-6
—— (*see* Moore et al). 1996;17:547-51
—— (*see* Wainwright et al). 1996;17:124-36
Herndon DN (*see* Mukhdomi et al). 1996;17:346-50
—— (*see* Zimmer et al). 1996;17:491-6
Hettich R (*see* Zwadlo-Klarwasser et al). 1996;17:287-93
Hickerson WL (*see* Smoot and Hickerson). 1996;17:273-9
Hobbs CL (*see* Harvey et al). 1996;17:176-80
Holaday M, Yarbrough A. Results of a hospital survey to determine the extent and type of psychologic services offered to patients with severe burns. 1996;17:280-4
Holguin PH, Rico AA, Garcia JP, Del Rio JL. Elbow anchylosis due to postburn heterotopic ossification. 1996;17:150-4
Hollender LG (*see* Fricke et al). 1996;17:338-45
Hughes HF (*see* Still et al). 1996;17:562-4

Hughes WB, DeClement FA, Hensell DO. Intradermal injection of epinephrine to decrease blood loss during split-thickness skin grafting. 1996;17:243-5
Hunt J. Development, implementation, and conclusions of critical pathways: walk don't run. 1996;17:S7-10
—— (*see* Grobmyer et al). 1996;17:532-9
—— (*see* Spann et al). 1996;17:515-7
—— (*see* Wainwright et al). 1996;17:124-36
Huribal M (*see* McMillen et al). 1996;17:384-9
Hurst TS (*see* Mosher et al). 1996;17:294-301
Hyland RH (*see* McIvor et al). 1996;17:237-40

I

Illmayer SJ (*see* Fraulin et al). 1996;17:19-29
Israeli A (*see* Ben-Bassat et al). 1996;17:421-8
Izenberg SD (*see* Frye et al). 1996;17:540-6

J

James LE (*see* Brown et al). 1996;17:552-7
Janus TJ, Barrash J. Neurologic and neurobehavioral effects of electric and lightning injuries. 1996;17:409-15
Jenkins HML, Blank V, Miller K, Turner J, Stanwick RS. A randomized single-blind evaluation of a discharge teaching book for pediatric patients with burns. 1996;17:49-61
Johnson DH (*see* Mosher et al). 1996;17:294-301
Johnson J (*see* Silverberg et al). 1996;17:252-9
—— (*see* Torres-Gray et al). 1996;17:161-8
Jones S (*see* Baker et al). 1996;17:327-33
Jönsson A (*see* Tarnow et al). 1996;17:207-12
Jordan MH (*see* Turner et al). 1996;17:169-74

K

Kagan R (*see* Wainwright et al). 1996;17:124-36
Kauhl W (*see* Zwadlo-Klarwasser et al). 1996;17:287-93
Kealey GP, Burger M. Critical pathways in clinical practice. 1996;17:S6-7
—— (*see* Foertsch et al). 1996;17:372-7
Keck T (*see* Richard et al). 1996;17:436-44
Keswani MH. The cost of burns and the relevance of prevention: the 1996 Everett Idris Evans Memorial Lecture. 1996;17:485-90
Kiessig V (*see* Rennekampff et al). 1996;17:213-21
—— (*see* Rennekampff et al). 1996;17:522-7
Kikuchi Y (*see* Klein et al). 1996;17:518-21
Kirby DL, O'Keefe JS, Neal JG, Bentrem DJ, Edlich RF. Does the architectural design of burn centers comply with the Americans with Disabilities Act? 1996;17:156-60
Klein GL, Kikuchi Y, Sherrard DJ, Simmons DJ, Biondo N, Traber DL. Burn-associated bone disease in sheep: roles of immobilization and endogenous corticosteroids. 1996;17:518-21
Klein MB, Shaw D, Barese S, Chapo GA, Cuono CB. A reliable and cost-effective in vitro assay of skin viability for skin banks and burn centers. 1996;17:565-70

L

LaBarbera MC (*see* Doong et al). 1996;17:497-514
LaLonde C, Nayak U, Hennigan J, Demling R. Antioxidants prevent the cellular deficit produced in response to burn injury. 1996;17:379-83
Landry VL. Animal extremists: a cause for concern? 1996;17:260-2
——. A software for power and statistical effect analysis. 1996;17:45-7
Law E (*see* Still et al). 1996;17:429-31 (Case rep.)
Law EJ (*see* Still et al). 1996;17:188-90
—— (*see* Still et al). 1996;17:562-4
Lawrence MH, de Riesthal HF, Calvano SE. Changes in memory and naive CD4+ lymphocytes in lymph nodes and spleen after thermal injury. 1996;17:1-6
LeDoux JM, Meyer WJ, Blakeney P, Herndon D. Positive self-regard as a coping mechanism for pediatric burn survivors. 1996;17:472-6
Lee RC (*see* Doong et al). 1996;17:497-513
Linder SA, Mele JA III, Harries T. Chronic hyperpigmentation from a heated mustard compress burn: a case report. 1996;17:351-2 (Case rep.)
Liu PY (*see* Orgill et al). 1996;17:311-22
Loomis W (*see* Rennekampff et al). 1996;17:522-7
Luterman A (*see* Frye et al). 1996;17:540-6
—— (*see* Wainwright et al). 1996;17:124-36
—— (*see* Watkins et al). 1996;17:78-92
Lynch JB (*see* Nanney et al). 1996;17:191-8

M

Madden M (*see* Wainwright et al). 1996;17:124-36
Maniscalco SP (*see* Grobmyer et al). 1996;17:532-9
Mantelle L (*see* Peck et al). 1996;17:39-44
Marshall GD (*see* Zimmer et al). 1996;17:491-6
Martin-Duffy K (*see* Baker et al). 1996;17:327-33
Marvin J. Closing remarks. 1996;17:S10-11
Mast B (*see* Bettinger et al). 1996;17:302-4
May SR (*see* Watkins et al). 1996;17:78-92
Mayers I (*see* Mosher et al). 1996;17:294-301
McCauley RL (*see* Mukhdomi et al). 1996;17:346-50
McDonald-Smith GP. NATIONAL TRACS™/ABA Burn Registry—the quest for quality burn data management. 1996;17:454-6
—— (*see* Miller and McDonald-Smith). 1996;17:369-70
McGlone JJ (*see* Penturf et al). 1996;17:465-70
McIvor RA, Zaidi J, Peters WJ, Hyland RH. Acute and chronic respiratory complications of toxic epidermal necrolysis. 1996;17:237-40
McLoughlin E (*see* Brigham and McLoughlin). 1996;17:95-107
McManus WF (*see* Harvey et al). 1996;17:176-80
McMillen MA, Huribal M, Cunningham ME, Bala RJ, Pleban WE, D'Aiuto ML. Endothelin-1, interleukin-6, and interleukin-8 levels increase in patients with burns. 1996;17:384-9
Mele JA III (*see* Linder et al). 1996;17:351-2 (Case rep.)
Merlo A (*see* Patiño et al). 1996;17:528-31

Mertz PM, Davis SC, Franzen L, Uchima F-D, Pickett MP, Pierschbacher MD, Polarek JW. Effects of an arginine–glycine–aspartic acid peptide-containing artificial matrix on epithelial migration in vitro and experimental second-degree burn wound healing in vivo. 1996;17:199-206
—— (*see* Davis et al). 1996;17:558-61
Meyer AA (*see* Tabata et al). 1996;17:231-6
Meyer W (*see* Moore et al). 1996;17:547-51
Meyer WJ (*see* LeDoux et al). 1996;17:472-6
Meyers R (*see* Stengle et al). 1996;17:14-8
Mileski WJ (*see* Spann et al). 1996;17:515-7
Miller K (*see* Jenkins et al). 1996;17:49-61
Miller S (*see* Hastings et al). 1996;17:416-20
—— (*see* Richard et al). 1996;17:444-53
Miller SF, McDonald-Smith GP. Report of the ABA meeting, March 13-17, 1996, Nashville, Tennessee. 1996;17:369-70
—— (*see* Staley et al). 1996;17:362-8
Minakawa H (*see* Yamamoto et al). 1996;17:397-401
Mitchell AT (*see* Mukhdomi et al). 1996;17:346-50
Mlakar J (*see* Torres-Gray et al). 1996;17:161-8
Moffat M (*see* Silverberg et al). 1996;17:252-9
Monafo W (*see* Wainwright et al). 1996;17:124-36
Mone M (*see* Saffle et al). 1996;17:353-61
Moore M (*see* Moore et al). 1996;17:547-51
Moore P, Moore M, Blakeney P, Meyer W, Murphy L, Herndon D. Competence and physical impairment of pediatric survivors of burns of more than 80% total body surface area. 1996;17:547-51
Morris SE (*see* Saffle et al). 1996;17:353-61
Mosher M, Hurst TS, Mayers I, Johnson DH. Lazaroids—not nitric oxide synthetase inhibitors—improve hemodynamics after thermal injury in anesthetized guinea pigs. 1996;17:294-301
Mozingo DW (*see* Harvey et al). 1996;17:176-80
Mukhdomi GJ, McCauley RL, Desai MH, Mitchell AT, Herndon DN. Cellulitis associated with burn scars: a retrospective review. 1996;17:346-50
Munster AM. Burns of the world. 1996;17:477-84
Murphy L (*see* Moore et al). 1996;17:547-51
Murphy M (*see* Punjabi et al). 1996;17:285 (Letter)

N

Nanney LB, Wenczak BA, Lynch JB. Progressive burn injury documented with vimentin immunostaining. 1996;17:191-8
Nayak U (*see* LaLonde et al). 1996;17:379-83
Neal JG (*see* Kirby et al). 1996;17:156-60
Nellgård P (*see* Tarnow et al). 1996;17:207-12
Nelson EW (*see* Winslow et al). 1996;17:117-23
Nolan D (*see* Hastings et al). 1996;17:416-20

O

O'Hara MW (*see* Foertsch et al). 1996;17:372-7
O'Keefe JS (*see* Kirby et al). 1996;17:156-60
Omnell L (*see* Fricke et al). 1996;17:338-45
Orgill DP, Liu PY, Ritterbush LS, Skrabut EM, Samuels JA, Shames SL. Debridement of porcine burns with a highly purified, ananain-based cysteine protease preparation. 1996;17:311-22
—— (*see* Abramson et al). 1996;17:402-8
Orlet HK (*see* Still et al). 1996;17:188-90

P

Parks DH (*see* Covington et al). 1996;17:222-30
Patiño O, Grana D, Bolgiani A, Prezzavento G, Merlo A, Mino J, Benaim F. Pulsed electromagnetic fields in experimental cutaneous wound healing in rats. 1996;17:528-31
Peck MD, Mantelle L, Ward CG. Comparison of length of hospital stay to mortality rate in a regional burn center. 1996;17:39-44
Peled IJ. Fluorescein and burn depth. 1996;17:378 (Letter)
Penturf ME, McGlone JJ, Griswold JA. Modulation of immune response in thermal injury by essential fatty acid–deficient diet. 1996;17:465-70
Peters WJ (*see* McIvor et al). 1996;17:237-40
Pickens HC (*see* Still et al). 1996;17:188-90
Pickett MP (*see* Mertz et al). 1996;17:199-206
Pierschbacher MD (*see* Mertz et al). 1996;17:199-206
Pleban WE (*see* McMillen et al). 1996;17:384-9
Polarek JW (*see* Cooper et al). 1996;17:108-16
—— (*see* Mertz et al). 1996;17:199-206
Pollard V (*see* Zimmer et al). 1996;17:491-6
Prezavento G (*see* Patiño et al). 1996;17:528-31
Pribaz JJ (*see* Abramson et al). 1996;17:402-8
Prough D (*see* Zimmer et al). 1996;17:491-6
Pruitt BA Jr (*see* Harvey et al). 1996;17:176-80
Punjabi AP, Bradrick JP, Murphy M. Stabilization of endotracheal tubes in children. 1996;17:285 (Letter)
Purdue G (*see* Spann et al). 1996;17:515-7
Purdue GF (*see* Grobmyer et al). 1996;17:532-9
Purvis RJ (*see* Watkins et al). 1996;17:78-92
Pyle J (*see* Stengle et al). 1996;17:14-8

R

Rennekampff HO, Hansbrough JF, Woods V Jr, Kiessig V. Integrin and matrix molecule expression in cultured skin replacements. 1996;17:213-21
——, Kiessig V, Loomis W, Hansbrough JF. Growth peptide release from biologic dressings: a comparison. 1996;17:522-7
Richard R (*see* Staley and Richard). 1996;17:S12-4
——, Staley M, Miller S, Warden G. To splint or not to splint—past philosophy and present practice: part I. 1996;17:444-53
——, Steinlage R, Staley M, Keck T. Mathematic model to estimate change in burn scar length required for joint range of motion. 1996;17:436-44
—— (*see* Staley et al). 1996;17:362-8
Rico AA (*see* Holguin et al). 1996;17:150-4

Ritterbush LS (*see* Orgill et al). 1996;17:311-22
Rodriguez DJ. Nutrition in patients with severe burns: state of the art. 1996;17:62-70
Ron M (*see* Ben-Bassat et al). 1996;17:421-8
Roth A (*see* Hastings et al). 1996;17:416-20

S

Sadinski C (*see* Baker et al). 1996;17:327-33
Saffle JR, Tuohig GM, Sullivan JJ, Shelby J, Morris SE, Mone M. Return to work as a measure of outcome in adults hospitalized for acute burn treatment. 1996;17:353-61
—— (*see* Winslow et al). 1996;17:117-23
Samuels JA (*see* Orgill et al). 1996;17:311-22
Sanders C (*see* Baker et al). 1996;17:327-33
Sanford S, Gore D. Unna's boot dressings facilitate outpatient skin grafting of hands. 1996;17:323-6
Scheirer R (*see* Still et al). 1996;17:429-31 (Case rep.)
Schiller WR (*see* Caruso et al). 1996;17:432-4 (Case rep.)
Schmitz C (*see* Zwadlo-Klarwasser et al). 1996;17:287-93
Shames SL (*see* Orgill et al). 1996;17:311-22
Shaw D (*see* Klein et al). 1996;17:565-70
Shelby J (*see* Saffle et al). 1996;17:353-61
—— (*see* Winslow et al). 1996;17:117-23
Sherrard DJ (*see* Klein et al). 1996;17:518-21
Shuster DB (*see* Staley et al). 1996;17:362-8
Silverberg R, Johnson J, Moffat M. The effects of soft tissue mobilization on the immature burn scar: results of a pilot study. 1996;17:252-9
Simmons DJ (*see* Klein et al). 1996;17:518-21
Simpson RL (*see* Yelon et al). 1996;17:241-2 (Case rep.)
Sittig K (*see* Wainwright et al). 1996;17:124-36
Sjöberg F, Gupta A. Anesthesia and thermal injury. 1996;17:285-6 (Letter)
Skrabut EM (*see* Orgill et al). 1996;17:311-22
Smoot EC III. Modified use of extremity tourniquets for burn wound debridement. 1996;17:334-7
——, Hickerson WL. Propane gas dangers and strategies for prevention of injuries. 1996;17:273-9
Spann K, Mileski WJ, Atiles L, Purdue G, Hunt J. Use of a pneumatonometer in burn scar assessment. 1996;17:515-7
Staley M, Richard R. Critical pathways to enhance the rehabilitation of patients with burns. 1996;17:S12-4
——, ——, Warden GD, Miller SF, Shuster DB. Functional outcomes for the patient with burn injuries. 1996;17:362-8
—— (*see* Richard et al). 1996;17:436-44
—— (*see* Richard et al). 1996;17:444-53
Stanwick RS (*see* Jenkins et al). 1996;17:49-61
Steinlage R (*see* Richard et al). 1996;17:436-44
Stengle J, Meyers R, Pyle J, Dries DJ. Neutrophil recruitment after remote scald injury. 1996;17:14-8
Still J, Scheirer R, Law E. Acute cholecystectomy performed through cultured epithelial autografts in a patient with burn injuries: a case report. 1996;17:429-31 (Case rep.)
Still JM, Law EJ, Duncan JW, Hughes HF. Long thoracic nerve injury due to an electric burn. 1996;17:562-4
——, Orlet HK, Law EJ, Pickens HC. Burns due to flammable solvents ignited with floor buffers. 1996;17:188-90
Still JM Jr (*see* Watkins et al). 1996;17:78-92
Stoddard FJ (*see* Foertsch et al). 1996;17:372-7
Strauss N (*see* Ben-Bassat et al). 1996;17:421-8
Sugihara T (*see* Yamamoto et al). 1996;17:397-401
Sullivan JJ (*see* Saffle et al). 1996;17:353-61
Sutherland S. Procedural burn pain intensity under conditions of varying physical control by the patient. 1996;17:457-63

T

Tabata T, deSerres S, Meyer AA. Differences in IgM synthesis to gut bacterial peptidoglycan polysaccharide after burn injury and gut ischemia. 1996;17:231-6
Tarnow P, Jönsson A, Nellgård P, Cassuto J. Reduced albumin extravasation in experimental rat skin and muscle burn injury by D-*myo*-inositol-1,2,6-triphosphate treatment. 1996;17:207-12
Torres-Gray D, Johnson J, Mlakar J. Rehabilitation of the burned hand: questionnaire results. 1996;17:161-8
Traber D (*see* Zimmer et al). 1996;17:491-6
Traber DL (*see* Klein et al). 1996;17:518-21
Tredget EE (*see* Fraulin et al). 1996;17:19-29
Trout S (*see* Dattolo et al). 1996;17:182-7
Tuohig GM (*see* Saffle et al). 1996;17:353-61
Turner DG, Berger N, Weiland AP, Jordan MH. The revised burn diagram and its effect on diagnosis-related group coding. 1996;17:169-74
Turner J (*see* Jenkins et al). 1996;17:49-61

U

Uchima F-D (*see* Mertz et al). 1996;17:199-206

V

Valentine S (*see* Baker et al). 1996;17:327-33

W

Wainwright D, Madden M, Luterman A, Hunt J, Monafo W, Heimbach D, Kagan R, Sittig K, Dimick A, Herndon D. Clinical evaluation of an acellular allograft dermal matrix in full-thickness burns. 1996;17:124-36
Wainwright DJ (*see* Covington et al). 1996;17:222-30
Wallace BH (*see* Caldwell et al). 1996;17:137-46
Ward CG (*see* Peck et al). 1996;17:39-44
Warden G (*see* Richard et al). 1996;17:444-53
Warden GD (*see* Brown et al). 1996;17:34-8
—— (*see* Brown et al). 1996;17:552-7
—— (*see* Staley et al). 1996;17:362-8
Washam MA (*see* Brown et al). 1996;17:552-7
Watkins PN, Cook EL, May SR, Still JM Jr, Luterman A, Purvis RJ. Postburn psychologic adaptation of family members of patients with burns. 1996;17:78-92

Weier HG (*see* Young et al). 1996;17:305-10
Weiland AP (*see* Turner et al). 1996;17:169-74
Wenczak BA (*see* Nanney et al). 1996;17:191-8
Wexler MR (*see* Ben-Bassat et al). 1996;17:421-8
Williams MD (*see* Frye et al). 1996;17:540-6
Winslow GA, Shelby J, Nelson EW, Saffle JR. Influence of allogeneic blood transfusion on natural killer cell activity in burn-injured mice. 1996;17:117-23
Woods V Jr (*see* Rennekampff et al). 1996;17:213-21

X

Xu W, Germain L, Goulet F, Auger FA. Permanent grafting of living skin substitutes: surgical parameters to control for successful results. 1996;17:7-13

Y

Yamamoto Y, Yokoyama T, Minakawa H, Sugihara T. Use of the expanded skin flap in esthetic reconstruction of postburn deformity. 1996;17:397-401
Yarbrough A (*see* Holaday and Yarbrough). 1996;17:280-4
Yarbrough DR III. Experience with toxic epidermal necrolysis treated in a burn center. 1996;17:30-3
Yelon JA, Simpson RL, Gudjonsson O. Formic acid inhalation injury: a case report. 1996;17:241-2 (Case rep.)
Yokoyama T (*see* Yamamoto et al). 1996;17:397-401
Young DM, Greulich KM, Weier HG. Species-specific in situ hybridization with fluorochrome-labeled DNA probes to study vascularization of human skin grafts on athymic mice. 1996;17:305-10
Yurko L, Fratianne RB. Clinical pathways as an education tool. 1996;17:S2-3

Z

Zaidi J (*see* McIvor et al). 1996;17:237-40
Zimmer S, Pollard V, Marshall GD, Garofalo RP, Traber D, Herndon DN, Prough D. Effects of endotoxin on the Th1/Th2 response in humans. 1996;17:491-6
Zwadlo-Klarwasser G, Kauhl W, Schmitz C, Hettich R. Influence of severe burn injury on the expression of RM 3/1 and HLA-DR antigens in human blood monocytes. 1996;17:287-93

Cumulative Subject Index, Volume 17*

A

Abnormalities
Use of the expanded skin flap in esthetic reconstruction of postburn deformity (Yamamoto, et al). 1996;17:397-401

Actins
Calcium antagonists alter cell shape and induce procollagenase synthesis in keloid and normal human dermal fibroblasts (Doong et al). 1996;17:497-514

Adaptation, psychological
Degree of burn, location of burn, and length of hospital stay as predictors of psychosocial status and physical functioning (Baker et al). 1996;17:327-33
Parent participation during burn debridement in relation to behavioral distress (Foertsch et al). 1996;17:372-7
Positive self-regard as a coping mechanism for pediatric burn survivors (LeDoux et al). 1996;17:472-6
Postburn psychologic adaptation of family members of patients with burns (Watkins et al). 1996;17:78-92
Results of a hospital survey to determine the extent and type of psychologic services offered to patients with severe burns (Holaday and Yarbrough). 1996;17:280-4

Adenine nucleotides
Antioxidants prevent the cellular deficit produced in response to burn injury (LaLonde et al). 1996;17:379-83

Adenosine triphosphate
Antioxidants prevent the cellular deficit produced in response to burn injury (LaLonde et al). 1996;17:379-83

Administration, topical
Sequential excision and grafting of the burn injuries of 1507 patients treated between 1967 and 1986: end results and the determinants of death (Caldwell et al). 1996;17:137-46

Adrenal cortex hormones
Burn-associated bone disease in sheep: roles of immobilization and endogenous corticosteroids (Klein et al). 1996;17:518-21

Aerosols
Formic acid inhalation injury: a case report (Yelon et al). 1996;17:241-2 (Case rep.)

Affect
Neurologic and neurobehavioral effects of electric and lightning injuries (Janus and Barrash). 1996;17:409-15

Aged
Prognostic indicators in the elderly patient with burns (Covington et al). 1996;17:222-30

Agitation
The use of haloperidol in the agitated, critically ill pediatric patient with burns (Brown et al). 1996;17:34-8

Albumins
Reduced albumin extravasation in experimental rat skin and muscle burn injury by D-*myo*-inositol-1,2,6-triphosphate treatment (Tarnow et al). 1996;17:207-12

Alcoholic intoxication
Alcohol, drug intoxication, or both at the time of burn injury as a predictor of complications and mortality in hospitalized patients with burns (Grobmyer et al). 1996;17:532-9

Alcoholism
Substance abuse in victims of fire (Barillo and Goode). 1996;17:71-6

Allografts; *see* **Transplantation, homologous**

American Burn Association
Report of the ABA meeting, March 13-17, 1996, Nashville, Tennessee (Miller and McDonald-Smith). 1996;17:369-70

Americans with Disabilities Act
Does the architectural design of burn centers comply with the Americans with Disabilities Act? (Kirby et al). 1996;17:156-60

Ananain-based, debriding enzymes
Debridement of porcine burns with a highly purified, ananain-based cysteine protease preparation (Orgill et al). 1996;17:311-22

Anesthesia
Anesthesia and thermal injury (Sjöberg and Gupta). 1996;17:285-6 (Letter)

Animal rights
Animal extremists: a cause for concern? (Landry). 1996;17:260-2

Ankylosis
Elbow anchylosis due to postburn heterotopic ossification (Holguin et al). 1996;17:150-4

Antibiotics
Sequential excision and grafting of the burn injuries of 1507 patients treated between 1967 and 1986: end results and the determinants of death (Caldwell et al). 1996;17:137-46

Antibody formation
Modulation of immune response in thermal injury by essential fatty acid–deficient diet (Penturf et al). 1996;17:465-70

Antigens
Influence of severe burn injury on the expression of RM 3/1 and HLA-DR antigens in human blood monocytes (Zwadlo-Klarwasser et al). 1996;17:287-93

*January/February, pp. 1-94; March/April, pp. 95-190; May/June, pp. 191-286; July/August, pp. 287-378; September/October, pp. 379-476; November/December, Part 1, 477-570. November/December, Part 2, pp. S1-S36.

Antioxidants
Antioxidants prevent the cellular deficit produced in response to burn injury (LaLonde et al). 1996;17: 379-83
Architecture
Does the architectural design of burn centers comply with the Americans with Disabilities Act? (Kirby et al). 1996;17:156-60
Arginine
The effect of an arginine–glycine–aspartic acid peptide and hyaluronate synthetic matrix on epithelialization of meshed skin graft interstices (Cooper et al). 1996; 17:108-16
Effects of an arginine–glycine–aspartic acid peptide-containing artificial matrix on epithelial migration in vitro and experimental second-degree burn wound healing in vivo (Mertz et al). 1996;17:199-206
Arm
Computer-assisted evaluation of hand and arm function after thermal injury (Harvey et al). 1996;17:176-80
Artificial intelligence
Simulated biologic intelligence used to predict length of stay and survival of burns (Frye et al). 1996;17:540-6
Aspartic acid
The effect of an arginine–glycine–aspartic acid peptide and hyaluronate synthetic matrix on epithelialization of meshed skin graft interstices (Cooper et al). 1996; 17:108-16
Effects of an arginine–glycine–aspartic acid peptide-containing artificial matrix on epithelial migration in vitro and experimental second-degree burn wound healing in vivo (Mertz et al). 1996;17:199-206
Autograft; *see* **Transplantation, autologous**

B

Bandages
Assessment of cosmetic and functional results of conservative versus surgical management of facial burns (Fraulin et al). 1996;17:19-29
Dermasorb versus Jelonet in patients with burns skin graft donor sites (Cadier and Clarke). 1996;17: 246-51
Proper use of nonadherent impregnated dressings (Britton). 1996;17:93-4 (Letter)
Skeletal and dental disturbances after facial burns and pressure garments (Fricke et al). 1996;17:338-45
Unna's boot dressings facilitate outpatient skin grafting of hands (Sanford and Gore). 1996;17:323-6
Wound coverage: Is there a difference? (Hastings et al). 1996;17:416-20
Basal metabolism
Nutrition in patients with severe burns: state of the art (Rodriguez). 1996;17:62-70
Baths
Procedural burn pain intensity under conditions of varying physical control by the patient (Sutherland). 1996;17:457-63
Behavior
Neurologic and neurobehavioral effects of electric and lightning injuries (Janus and Barrash). 1996;17:409-15
Biological assay
A reliable and cost-effective in vitro assay of skin viability for skin banks and burn centers (Klein et al). 1996; 17:565-70
Biological dressings
Experience with toxic epidermal necrolysis treated in a burn center (Yarbrough). 1996;17:30-3
Growth peptide release from biologic dressings: a comparison (Rennekampff et al). 1996;17:522-7
Blood loss, surgical
Intradermal injection of epinephrine to decrease blood loss during split-thickness skin grafting (Hughes et al). 1996;17:243-5
Modified use of extremity tourniquets for burn wound debridement (Smoot). 1996;17:334-7
Blood pressure
Anesthesia and thermal injury (Sjöberg and Gupta). 1996;17:285-6 (Letter)
Blood transfusion
Influence of allogeneic blood transfusion on natural killer cell activity in burn-injured mice (Winslow et al). 1996;17:117-23
Body surface area
Competence and physical impairment of pediatric survivors of burns of more than 80% total body surface area (Moore et al). 1996;17:547-51
The revised burn diagram and its effect on diagnosis-related group coding (Turner et al). 1996;17:169-74
Sequential excision and grafting of the burn injuries of 1507 patients treated between 1967 and 1986: end results and the determinants of death (Caldwell et al). 1996;17:137-46
The use of skin from a monozygotic twin combined with cultured epithelial autografts as coverage for a large surface area burn: a case report and review of the literature (Caruso et al). 1996;17:432-4 (Case rep.)
Bone development
Skeletal and dental disturbances after facial burns and pressure garments (Fricke et al). 1996;17:338-45
Bone diseases
Burn-associated bone disease in sheep: roles of immobilization and endogenous corticosteroids (Klein et al). 1996;17:518-21
Burn units
Comparison of length of hospital stay to mortality rate in a regional burn center (Peck et al). 1996;17:39-44
Does the architectural design of burn centers comply with the Americans with Disabilities Act? (Kirby et al). 1996;17:156-60
A reliable and cost-effective in vitro assay of skin viability for skin banks and burn centers (Klein et al). 1996; 17:565-70
Burns, chemical
Formic acid inhalation injury: a case report (Yelon et al). 1996;17:241-2 (Case rep.)
Burns, electric
Long thoracic nerve injury due to an electric burn (Still et al). 1996;17:562-4
Neurologic and neurobehavioral effects of electric and lightning injuries (Janus and Barrash). 1996;17:409-15

Persistent peripheral nerve pathologies in patients with electric burns (Haberal et al). 1996;17:147-9

Burns, inhalation

Formic acid inhalation injury: a case report (Yelon et al). 1996;17:241-2 (Case rep.)

Inhalation injury severity scoring system: a quantitative method (Brown et al). 1996;17:552-7

Burn-TRACS

Report of the ABA meeting, March 13-17, 1996, Nashville, Tennessee (Miller and McDonald-Smith). 1996; 17:369-70

C

Calcium channel blockers

Calcium antagonists alter cell shape and induce procollagenase synthesis in keloid and normal human dermal fibroblasts (Doong et al). 1996;17:497-514

Caloric intake

Nutrition in patients with severe burns: state of the art (Rodriguez). 1996;17:62-70

Carbohydrates

Nutrition in patients with severe burns: state of the art (Rodriguez). 1996;17:62-70

CD4-positive lymphocytes; *see* T4 lymphocytes

Cell size

Calcium antagonists alter cell shape and induce procollagenase synthesis in keloid and normal human dermal fibroblasts (Doong et al). 1996;17:497-513

Cellular immunity; *see* Immunity, cellular

Cellulitis

Cellulitis associated with burn scars: a retrospective review (Mukhdomi et al). 1996;17:346-50

Central nervous system

Persistent peripheral nerve pathologies in patients with electric burns (Haberal et al). 1996;17:147-9

Cephalometry

Skeletal and dental disturbances after facial burns and pressure garments (Fricke et al). 1996;17:338-45

Child

Competence and physical impairment of pediatric survivors of burns of more than 80% total body surface area (Moore et al). 1996;17:547-51

Parent participation during burn debridement in relation to behavioral distress (Foertsch et al). 1996;17: 372-7

Positive self-regard as a coping mechanism for pediatric burn survivors (LeDoux et al). 1996;17:472-6

A randomized single-blind evaluation of a discharge teaching book for pediatric patients with burns (Jenkins et al). 1996;17:49-61

Skeletal and dental disturbances after facial burns and pressure garments (Fricke et al). 1996;17:338-45

Stabilization of endotracheal tubes in children (Punjabi et al). 1996;17:285 (Letter)

The use of haloperidol in the agitated, critically ill pediatric patient with burns (Brown et al). 1996;17: 34-8

Cholecystectomy

Acute cholecystectomy performed through cultured epithelial autografts in a patient with burn injuries: a case report (Still et al). 1996;17:429-31 (Case rep.)

Cicatrix

Assessment of cosmetic and functional results of conservative versus surgical management of facial burns (Fraulin et al). 1996;17:19-29

Cellulitis associated with burn scars: a retrospective review (Mukhdomi et al). 1996;17:346-50

The effects of soft tissue mobilization on the immature burn scar: results of a pilot study (Silverberg et al). 1996;17:252-9

Hyaluronic acid impedes reepithelialization of skin graft donor sites (Bettinger et al). 1996;17:302-4

Mathematic model to estimate change in burn scar length required for joint range of motion (Richard et al). 1996;17:436-44

Use of a pneumatonometer in burn scar assessment (Spann et al). 1996;17:515-7

Cicatrix, hypertrophic

Chronic hyperpigmentation from a heated mustard compress burn: a case report (Linder et al). 1996;17: 351-2 (Case rep.)

Skeletal and dental disturbances after facial burns and pressure garments (Fricke et al). 1996;17:338-45

Clinical pathways

Clinical pathways as an education tool (Yurko and Fratianne). 1996;17:S2-3

Clinical practice

Critical pathways in clinical practice (Kealey and Burger). 1996;17:S6

Clinical Research Award

Use of a pneumatonometer in burn scar assessment (Spann et al). 1996;17:515-7

Coding; *see* Forms and records control

Cognition

Neurologic and neurobehavioral effects of electric and lightning injuries (Janus and Barrash). 1996;17:409-15

Collagen

Debridement of porcine burns with a highly purified, ananain-based cysteine protease preparation (Orgill et al). 1996;17:311-22

Collagenases

Calcium antagonists alter cell shape and induce procollagenase synthesis in keloid and normal human dermal fibroblasts (Doong et al). 1996;17:497-514

Colloids

Dermasorb versus Jelonet in patients with burns skin graft donor sites (Cadier and Clarke). 1996;17:246-51

Computer-assisted evaluation

Computer-assisted evaluation of hand and arm function after thermal injury (Harvey et al). 1996;17:176-80

Computers

NATIONAL TRACS™/ABA Burn Registry—the quest for quality burn data management (McDonald-Smith). 1996;17:454-6

Report of the ABA meeting, March 13-17, 1996, Nashville, Tennessee (Miller and McDonald-Smith). 1996; 17:369-70

A software for power and statistical effect analysis (Landry). 1996;17:45-7

Contracture
Mathematic model to estimate change in burn scar length required for joint range of motion (Richard et al). 1996;17:436-44
Coping; *see* **Adaptation, psychological**
Corticosteroids; *see* **Adrenal cortex hormones**
Costs and cost analysis
The cost of burns and the relevance of prevention: the 1996 Everett Idris Evans Memorial Lecture (Keswani). 1996;17:485-90
A reliable and cost-effective in vitro assay of skin viability for skin banks and burn centers (Klein et al). 1996; 17:565-70
Critical care
The use of haloperidol in the agitated, critically ill pediatric patient with burns (Brown et al). 1996;17:34-8
Critical pathways
Closing remarks (Marvin). 1996;17:S10-11
Critical pathways in clinical practice (Kealey and Burger). 1996;17:S6-7
Critical pathways in the QI process (Cornwell). 1996; 17:S3-6
Critical pathways to enhance the rehabilitation of patients with burns (Staley and Richard). 1996;17:S12-4
Development, implementation, and conclusions of critical pathways: walk don't run (Hunt). 1996;17:S7-10
Introduction (Gordon). 1996;17:S1
Cryopreservation
Growth peptide release from biologic dressings: a comparison (Rennekampff et al). 1996;17:522-7
Transplantation performance of human skin cryopreserved by programmed or stepwise freezing and stored at -80°C or -180°C (Ben-Bassat et al). 1996; 17:421-8
Cysteine proteinases
Debridement of porcine burns with a highly purified, ananain-based cysteine protease preparation (Orgill et al). 1996;17:311-22
Cytokines
Effects of endotoxin on the Th1/Th2 response in humans (Zimmer et al). 1996;17:491-6
Endothelin-1, interleukin-6, and interleukin-8 levels increase in patients with burns (McMillen et al). 1996; 17:384-9

D

Database management systems
NATIONAL TRACS™/ABA Burn Registry—the quest for quality burn data management (McDonald-Smith). 1996;17:454-6
Debridement
Assessment of cosmetic and functional results of conservative versus surgical management of facial burns (Fraulin et al). 1996;17:19-29
Debridement of porcine burns with a highly purified, ananain-based cysteine protease preparation (Orgill et al). 1996;17:311-22
Early debridement of second-degree burn wounds enhances the rate of epithelization—an animal model to evaluate burn wound therapies (Davis et al). 1996;17: 558-61
Modified use of extremity tourniquets for burn wound debridement (Smoot). 1996;17:334-7
Parent participation during burn debridement in relation to behavioral distress (Foertsch et al). 1996;17: 372-7
Deformities; *see* **Abnormalities**
Delirium
The use of haloperidol in the agitated, critically ill pediatric patient with burns (Brown et al). 1996;17:34-8
Depression
Results of a hospital survey to determine the extent and type of psychologic services offered to patients with severe burns (Holaday and Yarbrough). 1996; 17:280-4
Diagnosis-related groups
The revised burn diagram and its effect on diagnosis-related group coding (Turner et al). 1996;17:169-74
Diet
Modulation of immune response in thermal injury by essential fatty acid–deficient diet (Penturf et al). 1996; 17:465-70
Discharge; *see* **Patient discharge**
D-*myo*-inositol-1,2,6-triphosphate
Reduced albumin extravasation in experimental rat skin and muscle burn injury by D-*myo*-inositol-1,2,6-triphosphate treatment (Tarnow et al). 1996;17:207-12
DNA probes
Species-specific in situ hybridization with fluorochrome-labeled DNA probes to study vascularization of human skin grafts on athymic mice (Young et al). 1996; 17:305-10
Documentation
Functional outcomes for the patient with burn injuries (Staley et al). 1996;17:362-8
Progressive burn injury documented with vimentin immunostaining (Nanney et al). 1996;17:191-8
Dressings; *see* **Bandages**
Drug abuse; *see* **Substance abuse**

E

Ectropion
Assessment of cosmetic and functional results of conservative versus surgical management of facial burns (Fraulin et al). 1996;17:19-29
Edema
Reduced albumin extravasation in experimental rat skin and muscle burn injury by D-*myo*-inositol-1,2,6-triphosphate treatment (Tarnow et al). 1996;17:207-12
Education, continuing
Clinical pathways as an education tool (Yurko and Fratianne). 1996;17:S2-3
Elbow
Elbow anchylosis due to postburn heterotopic ossification (Holguin et al). 1996;17:150-4
Unna's boot dressings facilitate outpatient skin grafting of hands (Sanford and Gore). 1996;17:323-6
Elderly; *see* **Aged**
Electric injuries
Long thoracic nerve injury due to an electric burn (Still et al). 1996;17:562-4

Neurologic and neurobehavioral effects of electric and lightning injuries (Janus and Barrash). 1996;17:409-15
Persistent peripheral nerve pathologies in patients with electric burns (Haberal et al). 1996;17:147-9

Electromagnetics
Pulsed electromagnetic fields in experimental cutaneous wound healing in rats (Patiño et al). 1996;17:528-31

Endothelins
Endothelin-1, interleukin-6, and interleukin-8 levels increase in patients with burns (McMillen et al). 1996;17:384-9

Endotoxins
Effects of endotoxin on the Th1/Th2 response in humans (Zimmer et al). 1996;17:491-6

Energy metabolism
Antioxidants prevent the cellular deficit produced in response to burn injury (LaLonde et al). 1996;17:379-83

Enteral nutrition
Nutrition in patients with severe burns: state of the art (Rodriguez). 1996;17:62-70

Enzymes
Debridement of porcine burns with a highly purified, ananain-based cysteine protease preparation (Orgill et al). 1996;17:311-22

Epidermal necrolysis, toxic
Acute and chronic respiratory complications of toxic epidermal necrolysis (McIvor et al). 1996;17:237-40
Experience with toxic epidermal necrolysis treated in a burn center (Yarbrough). 1996;17:30-3

Epinephrine
Intradermal injection of epinephrine to decrease blood loss during split-thickness skin grafting (Hughes et al). 1996;17:243-5

Epithelium
Early debridement of second-degree burn wounds enhances the rate of epithelization—an animal model to evaluate burn wound therapies (Davis et al). 1996;17:558-61
Growth peptide release from biologic dressings: a comparison (Rennekampff et al). 1996;17:522-7

Essential fatty acids; *see* **Fatty acids, essential**

Esthetics
Assessment of cosmetic and functional results of conservative versus surgical management of facial burns (Fraulin et al). 1996;17:19-29
Use of the expanded skin flap in esthetic reconstruction of postburn deformity (Yamamoto et al). 1996;17:397-401

Everett Idris Evans Memorial Lecture
The cost of burns and the relevance of prevention: the 1996 Everett Idris Evans Memorial Lecture (Keswani). 1996;17:485-90

Exfoliative skin disorders
Experience with toxic epidermal necrolysis treated in a burn center (Yarbrough). 1996;17:30-3

Extracellular matrix
Clinical evaluation of an acellular allograft dermal matrix in full-thickness burns (Wainwright et al). 1996;17:124-36
The effect of an arginine–glycine–aspartic acid peptide and hyaluronate synthetic matrix on epithelialization of meshed skin graft interstices (Cooper et al). 1996;17:108-16
Effects of an arginine–glycine–aspartic acid peptide-containing artificial matrix on epithelial migration in vitro and experimental second-degree burn wound healing in vivo (Mertz et al). 1996;17:199-206
Integrin and matrix molecule expression in cultured skin replacements (Rennekampff et al). 1996;17:213-21

F

Face
Assessment of cosmetic and functional results of conservative versus surgical management of facial burns (Fraulin et al). 1996;17:19-29
Formic acid inhalation injury: a case report (Yelon et al). 1996;17:241-2 (Case rep.)
Skeletal and dental disturbances after facial burns and pressure garments (Fricke et al). 1996;17:338-45

Family
Postburn psychologic adaptation of family members of patients with burns (Watkins et al). 1996;17:78-92

Fats
Nutrition in patients with severe burns: state of the art (Rodriguez). 1996;17:62-70

Fatty acids, essential
Modulation of immune response in thermal injury by essential fatty acid–deficient diet (Penturf et al). 1996;17:465-70

Fibroblasts
Calcium antagonists alter cell shape and induce procollagenase synthesis in keloid and normal human dermal fibroblasts (Doong et al). 1996;17:497-514
The effect of an arginine–glycine–aspartic acid peptide and hyaluronate synthetic matrix on epithelialization of meshed skin graft interstices (Cooper et al). 1996;17:108-16
Effects of an arginine–glycine–aspartic acid peptide-containing artificial matrix on epithelial migration in vitro and experimental second-degree burn wound healing in vivo (Mertz et al). 1996;17:199-206

Fibronectins
The effect of an arginine–glycine–aspartic acid peptide and hyaluronate synthetic matrix on epithelialization of meshed skin graft interstices (Cooper et al). 1996;17:108-16
Effects of an arginine–glycine–aspartic acid peptide-containing artificial matrix on epithelial migration in vitro and experimental second-degree burn wound healing in vivo (Mertz et al). 1996;17:199-206
Integrin and matrix molecule expression in cultured skin replacements (Rennekampff et al). 1996;17:213-21

Fluid therapy
Experience with toxic epidermal necrolysis treated in a burn center (Yarbrough). 1996;17:30-3

Fluoresceins
Fluorescein and burn depth (Peled). 1996;17:378 (Letter)

Fluorescent dyes
Species-specific in situ hybridization with fluorochrome-labeled DNA probes to study vascularization of hu-

Fluorescent dyes—cont'd
man skin grafts on athymic mice (Young et al). 1996;17:305-10
Formic acids
Formic acid inhalation injury: a case report (Yelon et al). 1996;17:241-2 (Case rep.)
Forms and records control
The revised burn diagram and its effect on diagnosis-related group coding (Turner et al). 1996;17:169-74
Functional outcome report
Functional outcomes for the patient with burn injuries (Staley et al). 1996;17:362-8

G

Gases
Propane gas dangers and strategies for prevention of injuries (Smoot and Hickerson). 1996;17:273-9
Gasoline
Burns due to flammable solvents ignited with floor buffers (Still et al). 1996;17:188-90
Gelatin
Debridement of porcine burns with a highly purified, ananain-based cysteine protease preparation (Orgill et al). 1996;17:311-22
Glycine
The effect of an arginine–glycine–aspartic acid peptide and hyaluronate synthetic matrix on epithelialization of meshed skin graft interstices (Cooper et al). 1996;17:108-16
Effects of an arginine–glycine–aspartic acid peptide-containing artificial matrix on epithelial migration in vitro and experimental second-degree burn wound healing in vivo (Mertz et al). 1996;17:199-206
Graft survival
Permanent grafting of living skin substitutes: surgical parameters to control for successful results (Xu et al). 1996;17:7-13
Grafting, skin; *see* **Skin transplantation**
Grip strength
Computer-assisted evaluation of hand and arm function after thermal injury (Harvey et al). 1996;17:176-80
Growth substances
Effects of an arginine–glycine–aspartic acid peptide-containing artificial matrix on epithelial migration in vitro and experimental second-degree burn wound healing in vivo (Mertz et al). 1996;17:199-206
Growth peptide release from biologic dressings: a comparison (Rennekampff et al). 1996;17:522-7

H

Haloperidol
The use of haloperidol in the agitated, critically ill pediatric patient with burns (Brown et al). 1996;17:34-8
Hand
Computer-assisted evaluation of hand and arm function after thermal injury (Harvey et al). 1996;17:176-80
Rehabilitation of the burned hand: questionnaire results (Torres-Gray et al). 1996;17:161-8
Unna's boot dressings facilitate outpatient skin grafting of hands (Sanford and Gore). 1996;17:323-6
Health plan implementation
Development, implementation, and conclusions of critical pathways: walk don't run (Hunt). 1996;17:S7-10
Hemodynamics
Lazaroids—not nitric oxide synthetase inhibitors—improve hemodynamics after thermal injury in anesthetized guinea pigs (Mosher et al). 1996;17:294-301
Heterotopic ossification; *see* **Ossification, heterotopic**
HLA-DR antigens
Influence of severe burn injury on the expression of RM 3/1 and HLA-DR antigens in human blood monocytes (Zwadlo-Klarwasser et al). 1996;17:287-93
Home care services
Home health care and burn care: an educational and economical program (Dattolo et al). 1996;17:182-7
Hospital design and construction
Does the architectural design of burn centers comply with the Americans with Disabilities Act? (Kirby et al). 1996;17:156-60
Hospitalization
Burn incidence and medical care use in the United States: estimates, trends, and data sources (Brigham and McLoughlin). 1996;17:95-107
Humoral immunity; *see* **Antibody formation**
Hyaluronic acid
The effect of an arginine–glycine–aspartic acid peptide and hyaluronate synthetic matrix on epithelialization of meshed skin graft interstices (Cooper et al). 1996;17:108-16
Effects of an arginine–glycine–aspartic acid peptide-containing artificial matrix on epithelial migration in vitro and experimental second-degree burn wound healing in vivo (Mertz et al). 1996;17:199-206
Hyaluronic acid impedes reepithelialization of skin graft donor sites (Bettinger et al). 1996;17:302-4
Hydrotherapy
Clarification on the use of silver-based purification systems (Edlich). 1996;17:285 (Letter)
Hyperemia
Modified use of extremity tourniquets for burn wound debridement (Smoot). 1996;17:334-7
Hyperpigmentation
Chronic hyperpigmentation from a heated mustard compress burn: a case report (Linder et al). 1996;17:351-2 (Case rep.)
Hypnotics and sedatives
The use of haloperidol in the agitated, critically ill pediatric patient with burns (Brown et al). 1996;17:34-8

I

IgM
Differences in IgM synthesis to gut bacterial peptidoglycan polysaccharide after burn injury and gut ischemia (Tabata et al). 1996;17:231-6
Immobilization
Burn-associated bone disease in sheep: roles of immobilization and endogenous corticosteroids (Klein et al). 1996;17:518-21

Immune system
Essential microminerals and their response to burn injury (Gamliel et al). 1996;17:264-72
Modulation of immune response in thermal injury by essential fatty acid–deficient diet (Penturf et al). 1996; 17:465-70
Immunity, cellular
Modulation of immune response in thermal injury by essential fatty acid–deficient diet (Penturf et al). 1996; 17:465-70
Immunohistochemistry
Progressive burn injury documented with vimentin immunostaining (Nanney et al). 1996;17:191-8
Immunosuppression
Influence of severe burn injury on the expression of RM 3/1 and HLA-DR antigens in human blood monocytes (Zwadlo-Klarwasser et al). 1996;17:287-93
In situ hybridization
Species-specific in situ hybridization with fluorochrome-labeled DNA probes to study vascularization of human skin grafts on athymic mice (Young et al). 1996; 17:305-10
In vitro
A reliable and cost-effective in vitro assay of skin viability for skin banks and burn centers (Klein et al). 1996; 17:565-70
Infection
Cellulitis associated with burn scars: a retrospective review (Mukhdomi et al). 1996;17:346-50
Inflammation
Antioxidants prevent the cellular deficit produced in response to burn injury (LaLonde et al). 1996;17: 379-83
Reduced albumin extravasation in experimental rat skin and muscle burn injury by D-*myo*-inositol-1,2,6-tri-phosphate treatment (Tarnow et al). 1996;17: 207-12
Inhalation injury; *see* **Smoke inhalation injury**
Injections, intradermal
Intradermal injection of epinephrine to decrease blood loss during split-thickness skin grafting (Hughes et al). 1996;17:243-5
Injury severity score
Inhalation injury severity scoring system: a quantitative method (Brown et al). 1996;17:552-7
Integrins
Integrin and matrix molecule expression in cultured skin replacements (Rennekampff et al). 1996;17:213-21
Intelligence
Simulated biologic intelligence used to predict length of stay and survival of burns (Frye et al). 1996;17:540-6
Interferons
Effects of endotoxin on the Th1/Th2 response in humans (Zimmer et al). 1996;17:491-6
Interleukins
Effects of endotoxin on the Th1/Th2 response in humans (Zimmer et al). 1996;17:491-6
Endothelin-1, interleukin-6, and interleukin-8 levels increase in patients with burns (McMillen et al). 1996; 17:384-9
Intestines
Differences in IgM synthesis to gut bacterial peptidoglycan polysaccharide after burn injury and gut ischemia (Tabata et al). 1996;17:231-6
Intoxication; *see* **Poisoning**
Intubation
Stabilization of endotracheal tubes in children (Punjabi et al). 1996;17:285 (Letter)
Ischemia
Differences in IgM synthesis to gut bacterial peptidoglycan polysaccharide after burn injury and gut ischemia (Tabata et al). 1996;17:231-6

K

Keloid
Calcium antagonists alter cell shape and induce procollagenase synthesis in keloid and normal human dermal fibroblasts (Doong et al). 1996;17:497-513
Keratinocytes
The effect of an arginine–glycine–aspartic acid peptide and hyaluronate synthetic matrix on epithelialization of meshed skin graft interstices (Cooper et al). 1996; 17:108-16
Effects of an arginine–glycine–aspartic acid peptide-containing artificial matrix on epithelial migration in vitro and experimental second-degree burn wound healing in vivo (Mertz et al). 1996;17:199-206
Integrin and matrix molecule expression in cultured skin replacements (Rennekampff et al). 1996;17:213-21
Kerosene
Burns due to flammable solvents ignited with floor buffers (Still et al). 1996;17:188-90
Killer cells, natural
Influence of allogeneic blood transfusion on natural killer cell activity in burn-injured mice (Winslow et al). 1996;17:117-23
Modulation of immune response in thermal injury by essential fatty acid–deficient diet (Penturf et al). 1996; 17:465-70

L

Lazaroids
Lazaroids—not nitric oxide synthetase inhibitors—improve hemodynamics after thermal injury in anesthetized guinea pigs (Mosher et al). 1996;17:294-301
Length of stay
Comparison of length of hospital stay to mortality rate in a regional burn center (Peck et al). 1996;17:39-44
Degree of burn, location of burn, and length of hospital stay as predictors of psychosocial status and physical functioning (Baker et al). 1996;17:327-33
Simulated biologic intelligence used to predict length of stay and survival of burns (Frye et al). 1996;17:540-6
Letters
Letters to the editor. 285-6, 378, 1996;17:93-4
Lightning injuries
Neurologic and neurobehavioral effects of electric and lightning injuries (Janus and Barrash). 1996;17:409-15

Lindberg Award
Calcium antagonists alter cell shape and induce procollagenase synthesis in keloid and normal human dermal fibroblasts (Doong et al). 1996;17:497-514
Lipid peroxidation
Lazaroids—not nitric oxide synthetase inhibitors—improve hemodynamics after thermal injury in anesthetized guinea pigs (Mosher et al). 1996;17:294-301
Literature
Burns of the world (Munster). 1996;17:477-84
Lund and Browder diagram
The revised burn diagram and its effect on diagnosis-related group coding (Turner et al). 1996;17:169-74
Lung
Neutrophil recruitment after remote scald injury (Stengle et al). 1996;17:14-8
Lymph nodes
Changes in memory and naive CD4+ lymphocytes in lymph nodes and spleen after thermal injury (Lawrence et al). 1996;17:1-6

M

Mandible
Skeletal and dental disturbances after facial burns and pressure garments (Fricke et al). 1996;17:338-45
Massage
The effects of soft tissue mobilization on the immature burn scar: results of a pilot study (Silverberg et al). 1996;17:252-9
Mathematical model; *see* **Models, theoretical**
Maxilla
Skeletal and dental disturbances after facial burns and pressure garments (Fricke et al). 1996;17:338-45
Metabolism
Essential microminerals and their response to burn injury (Gamliel et al). 1996;17:264-72
Nutrition in patients with severe burns: state of the art (Rodriguez). 1996;17:62-70
Methohexital
Anesthesia and thermal injury (Sjöberg and Gupta). 1996;17:285-6 (Letter)
Microstomia
Assessment of cosmetic and functional results of conservative versus surgical management of facial burns (Fraulin et al). 1996;17:19-29
Minerals
Essential microminerals and their response to burn injury (Gamliel et al). 1996;17:264-72
Models, theoretical
Mathematic model to estimate change in burn scar length required for joint range of motion (Richard et al). 1996;17:436-44
Monocytes
Effects of endotoxin on the Th1/Th2 response in humans (Zimmer et al). 1996;17:491-6
Endothelin-1, interleukin-6, and interleukin-8 levels increase in patients with burns (McMillen et al). 1996;17:384-9
Influence of severe burn injury on the expression of RM 3/1 and HLA-DR antigens in human blood monocytes (Zwadlo-Klarwasser et al). 1996;17:287-93
Mood; *see* **Affect**
Mortality
Alcohol, drug intoxication, or both at the time of burn injury as a predictor of complications and mortality in hospitalized patients with burns (Grobmyer et al). 1996;17:532-9
Comparison of length of hospital stay to mortality rate in a regional burn center (Peck et al). 1996;17:39-44
Motor neurons
Persistent peripheral nerve pathologies in patients with electric burns (Haberal et al). 1996;17:147-9
Moyer Award
Effects of endotoxin on the Th1/Th2 response in humans (Zimmer et al). 1996;17:491-6
Mucous membrane
Acute and chronic respiratory complications of toxic epidermal necrolysis (McIvor et al). 1996;17:237-40
Muscles
Reduced albumin extravasation in experimental rat skin and muscle burn injury by D-*myo*-inositol-1,2,6-triphosphate treatment (Tarnow et al). 1996;17:207-12

N

NATIONAL TRACS™/ABA Burn Registry
NATIONAL TRACS™/ABA Burn Registry—the quest for quality burn data management (McDonald-Smith). 1996;17:454-6
Neovascularization
Substance P has a role in neurogenic mediation of human burn wound healing (Dunnick et al). 1996;17:390-6
Neurologic examination
Neurologic and neurobehavioral effects of electric and lightning injuries (Janus and Barrash). 1996;17:409-15
Neuropeptides
Substance P has a role in neurogenic mediation of human burn wound healing (Dunnick et al). 1996;17:390-6
Neutrophils
Neutrophil recruitment after remote scald injury (Stengle et al). 1996;17:14-8
Nitric oxide
Lazaroids—not nitric oxide synthetase inhibitors—improve hemodynamics after thermal injury in anesthetized guinea pigs (Mosher et al). 1996;17:294-301
Nucleotides
Antioxidants prevent the cellular deficit produced in response to burn injury (LaLonde et al). 1996;17:379-83
Nutrition
Essential microminerals and their response to burn injury (Gamliel et al). 1996;17:264-72
Experience with toxic epidermal necrolysis treated in a burn center (Yarbrough). 1996;17:30-3
Modulation of immune response in thermal injury by essential fatty acid–deficient diet (Penturf et al). 1996;17:465-70
Nutrition in patients with severe burns: state of the art (Rodriguez). 1996;17:62-70

O

Occlusive dressings
Sequential excision and grafting of the burn injuries of 1507 patients treated between 1967 and 1986: end results and the determinants of death (Caldwell et al). 1996;17:137-46
Occupational therapy
Functional outcomes for the patient with burn injuries (Staley et al). 1996;17:362-8
A randomized single-blind evaluation of a discharge teaching book for pediatric patients with burns (Jenkins et al). 1996;17:49-61
Rehabilitation of the burned hand: questionnaire results (Torres-Gray et al). 1996;17:161-8
Ossification, heterotopic
Elbow anchylosis due to postburn heterotopic ossification (Holguin et al). 1996;17:150-4
Outcome assessment
Functional outcomes for the patient with burn injuries (Staley et al). 1996;17:362-8
Return to work as a measure of outcome in adults hospitalized for acute burn treatment (Saffle et al). 1996;17:353-61

P

Pain
Procedural burn pain intensity under conditions of varying physical control by the patient (Sutherland). 1996;17:457-63
Parenteral nutrition
Nutrition in patients with severe burns: state of the art (Rodriguez). 1996;17:62-70
Parents
Parent participation during burn debridement in relation to behavioral distress (Foertsch et al). 1996;17:372-7
Patient discharge
Degree of burn, location of burn, and length of hospital stay as predictors of psychosocial status and physical functioning (Baker et al). 1996;17:327-33
A randomized single-blind evaluation of a discharge teaching book for pediatric patients with burns (Jenkins et al). 1996;17:49-61
Pentobarbital
Anesthesia and thermal injury (Sjöberg and Gupta). 1996;17:285-6 (Letter)
Peptides
The effect of an arginine–glycine–aspartic acid peptide and hyaluronate synthetic matrix on epithelialization of meshed skin graft interstices (Cooper et al). 1996;17:108-16
Effects of an arginine–glycine–aspartic acid peptide-containing artificial matrix on epithelial migration in vitro and experimental second-degree burn wound healing in vivo (Mertz et al). 1996;17:199-206
Growth peptide release from biologic dressings: a comparison (Rennekampff et al). 1996;17:522-7
Peptidoglycan
Differences in IgM synthesis to gut bacterial peptidoglycan polysaccharide after burn injury and gut ischemia (Tabata et al). 1996;17:231-6
Peripheral nervous system
Persistent peripheral nerve pathologies in patients with electric burns (Haberal et al). 1996;17:147-9
Petroleum
Propane gas dangers and strategies for prevention of injuries (Smoot and Hickerson). 1996;17:273-9
Physical therapy
The effects of soft tissue mobilization on the immature burn scar: results of a pilot study (Silverberg et al). 1996;17:252-9
Functional outcomes for the patient with burn injuries (Staley et al). 1996;17:362-8
A randomized single-blind evaluation of a discharge teaching book for pediatric patients with burns (Jenkins et al). 1996;17:49-61
Rehabilitation of the burned hand: questionnaire results (Torres-Gray et al). 1996;17:161-8
To splint or not to splint—past philosophy and present practice: part I (Richard et al). 1996;17:444-53
Pneumatonometer
Use of a pneumatonometer in burn scar assessment (Spann et al). 1996;17:515-7
Poisoning
Alcohol, drug intoxication, or both at the time of burn injury as a predictor of complications and mortality in hospitalized patients with burns (Grobmyer et al). 1996;17:532-9
Polysaccharides
Differences in IgM synthesis to gut bacterial peptidoglycan polysaccharide after burn injury and gut ischemia (Tabata et al). 1996;17:231-6
Polyurethanes
Dermasorb versus Jelonet in patients with burns skin graft donor sites (Cadier and Clarke). 1996;17:246-51
Porcine xenograft dressings; *see* **Biological dressings**
Presidential address
Burns of the world (Munster). 1996;17:477-84
Pressure
The effects of soft tissue mobilization on the immature burn scar: results of a pilot study (Silverberg et al). 1996;17:252-9
Pressure dressing
Assessment of cosmetic and functional results of conservative versus surgical management of facial burns (Fraulin et al). 1996;17:19-29
Skeletal and dental disturbances after facial burns and pressure garments (Fricke et al). 1996;17:338-45
Prevention and control
Burns due to flammable solvents ignited with floor buffers (Still et al). 1996;17:188-90
The cost of burns and the relevance of prevention: the 1996 Everett Idris Evans Memorial Lecture (Keswani). 1996;17:485-90
Propane gas dangers and strategies for prevention of injuries (Smoot and Hickerson). 1996;17:273-9
Substance abuse in victims of fire (Barillo and Goode). 1996;17:71-6
Procollagen
Calcium antagonists alter cell shape and induce procollagenase synthesis in keloid and normal human dermal fibroblasts (Doong et al). 1996;17:497-514

Program development
Development, implementation, and conclusions of critical pathways: walk don't run (Hunt). 1996;17:S7-10
Propane
Propane gas dangers and strategies for prevention of injuries (Smoot and Hickerson). 1996;17:273-9
Prospective payment system
Home health care and burn care: an educational and economical program (Dattolo et al). 1996;17:182-7
The revised burn diagram and its effect on diagnosis-related group coding (Turner et al). 1996;17:169-74
Proteins
Nutrition in patients with severe burns: state of the art (Rodriguez). 1996;17:62-70
Psychological tests
Results of a hospital survey to determine the extent and type of psychologic services offered to patients with severe burns (Holaday and Yarbrough). 1996;17:280-4
Pulmonary function tests; *see* **Respiratory function tests**

Q

Quality assurance
Critical pathways in the QI process (Cornwell). 1996;17:S4-5
Questionnaires
Rehabilitation of the burned hand: questionnaire results (Torres-Gray et al). 1996;17:161-8

R

Radiation
Pulsed electromagnetic fields in experimental cutaneous wound healing in rats (Patiño et al). 1996;17:528-31
Range of motion, articular
The effects of soft tissue mobilization on the immature burn scar: results of a pilot study (Silverberg et al). 1996;17:252-9
Mathematic model to estimate change in burn scar length required for joint range of motion (Richard et al). 1996;17:436-44
Registries
NATIONAL TRACS™/ABA Burn Registry—the quest for quality burn data management (McDonald-Smith). 1996;17.454-6
Rehabilitation
Critical pathways to enhance the rehabilitation of patients with burns (Staley and Richard). 1996;17:S12-4
Reimbursement mechanisms
Home health care and burn care: an educational and economical program (Dattolo et al). 1996;17:182-7
Research
Animal extremists: a cause for concern? (Landry). 1996;17:260-2
Use of a pneumatonometer in burn scar assessment (Spann et al). 1996;17:515-7
Respiration, artificial
Acute and chronic respiratory complications of toxic epidermal necrolysis (McIvor et al). 1996;17:237-40
Respiratory function tests
Acute and chronic respiratory complications of toxic epidermal necrolysis (McIvor et al). 1996;17:237-40
Return to work
Return to work as a measure of outcome in adults hospitalized for acute burn treatment (Saffle et al). 1996;17:353-61

S

Scars; *see* **Cicatrix**
Sedatives; *see* **Hypnotics and sedatives**
Self concept
Positive self-regard as a coping mechanism for pediatric burn survivors (LeDoux et al). 1996;17:472-6
Skin transplantation
Acute cholecystectomy performed through cultured epithelial autografts in a patient with burn injuries: a case report (Still et al). 1996;17:429-31 (Case rep.)
Assessment of cosmetic and functional results of conservative versus surgical management of facial burns (Fraulin et al). 1996;17:19-29
Clinical evaluation of an acellular allograft dermal matrix in full-thickness burns (Wainwright et al). 1996;17:124-36
The effect of an arginine–glycine–aspartic acid peptide and hyaluronate synthetic matrix on epithelialization of meshed skin graft interstices (Cooper et al). 1996;17:108-16
Growth peptide release from biologic dressings: a comparison (Rennekampff et al). 1996;17:522-7
Integrin and matrix molecule expression in cultured skin replacements (Rennekampff et al). 1996;17:213-21
Intradermal injection of epinephrine to decrease blood loss during split-thickness skin grafting (Hughes et al). 1996;17:243-5
Permanent grafting of living skin substitutes: surgical parameters to control for successful results (Xu et al). 1996;17:7-13
A reliable and cost-effective in vitro assay of skin viability for skin banks and burn centers (Klein et al). 1996;17:565-70
Sequential excision and grafting of the burn injuries of 1507 patients treated between 1967 and 1986: end results and the determinants of death (Caldwell et al). 1996;17:137-46
Species-specific in situ hybridization with fluorochrome-labeled DNA probes to study vascularization of human skin grafts on athymic mice (Young et al). 1996;17:305-10
Transplantation performance of human skin cryopreserved by programmed or stepwise freezing and stored at -80°C or -180°C (Ben-Bassat et al). 1996;17:421-8
Unna's boot dressings facilitate outpatient skin grafting of hands (Sanford and Gore). 1996;17:323-6
The use of skin from a monozygotic twin combined with cultured epithelial autografts as coverage for a large surface area burn: a case report and review of the literature (Caruso et al). 1996;17:432-4 (Case rep.)
Use of the expanded skin flap in esthetic reconstruction of postburn deformity (Yamamoto et al). 1996;17:397-401

Smoke inhalation injury
Inhalation injury severity scoring system: a quantitative method (Brown et al). 1996;17:552-7
Soft tissue mobilization
The effects of soft tissue mobilization on the immature burn scar: results of a pilot study (Silverberg et al). 1996;17:252-9
Software
A software for power and statistical effect analysis (Landry). 1996;17:45-7
Solvents
Burns due to flammable solvents ignited with floor buffers (Still et al). 1996;17:188-90
Spleen
Changes in memory and naive CD4+ lymphocytes in lymph nodes and spleen after thermal injury (Lawrence et al). 1996;17:1-6
Splints
To splint or not to splint—past philosophy and present practice: part I (Richard et al). 1996;17:444-53
Unna's boot dressings facilitate outpatient skin grafting of hands (Sanford and Gore). 1996;17:323-6
Stains and staining
Integrin and matrix molecule expression in cultured skin replacements (Rennekampff et al). 1996;17:213-21
Progressive burn injury documented with vimentin immunostaining (Nanney et al). 1996;17:191-8
Statistics
Burn incidence and medical care use in the United States: estimates, trends, and data sources (Brigham and McLoughlin). 1996;17:95-107
A software for power and statistical effect analysis (Landry). 1996;17:45-7
Stress disorders, post-traumatic
Results of a hospital survey to determine the extent and type of psychologic services offered to patients with severe burns (Holaday and Yarbrough). 1996;17:280-4
Substance abuse
Alcohol, drug intoxication, or both at the time of burn injury as a predictor of complications and mortality in hospitalized patients with burns (Grobmyer et al). 1996;17:532-9
Substance abuse in victims of fire (Barillo and Goode). 1996;17:71-6
Substance P
Substance P has a role in neurogenic mediation of human burn wound healing (Dunnick et al). 1996;17:390-6
Surgical flaps
Use of the expanded skin flap in esthetic reconstruction of postburn deformity (Yamamoto et al). 1996;17:397-401
Survival
Competence and physical impairment of pediatric survivors of burns of more than 80% total body surface area (Moore et al). 1996;17:547-51
Simulated biologic intelligence used to predict length of stay and survival of burns (Frye et al). 1996;17:540-6

T

T4 lymphocytes
Changes in memory and naive CD4+ lymphocytes in lymph nodes and spleen after thermal injury (Lawrence et al). 1996;17:1-6
Thoracic nerves
Long thoracic nerve injury due to an electric burn (Still et al). 1996;17:562-4
Tissue donors
The use of skin from a monozygotic twin combined with cultured epithelial autografts as coverage for a large surface area burn: a case report and review of the literature (Caruso et al). 1996;17:432-4 (Case rep.)
Tissue transplantation
The use of free tissue transfer in burn reconstruction (Abramson et al). 1996;17:402-8
T-lymphocytes
Influence of severe burn injury on the expression of RM 3/1 and HLA-DR antigens in human blood monocytes (Zwadlo-Klarwasser et al). 1996;17:287-93
Modulation of immune response in thermal injury by essential fatty acid–deficient diet (Penturf et al). 1996;17:465-70
T-lymphocytes, helper-induced
Effects of endotoxin on the Th1/Th2 response in humans (Zimmer et al). 1996;17:491-6
Tooth
Skeletal and dental disturbances after facial burns and pressure garments (Fricke et al). 1996;17:338-45
Total active motion
Computer-assisted evaluation of hand and arm function after thermal injury (Harvey et al). 1996;17:176-80
Total quality management
Critical pathways in the QI process (Cornwell). 1996;17:S4-5
Tourniquets
Modified use of extremity tourniquets for burn wound debridement (Smoot). 1996;17:334-7
Trace elements
Essential microminerals and their response to burn injury (Gamliel et al). 1996;17:264-72
Transplantation, autologous
Acute cholecystectomy performed through cultured epithelial autografts in a patient with burn injuries: a case report (Still et al). 1996;17:429-31 (Case rep.)
Permanent grafting of living skin substitutes: surgical parameters to control for successful results (Xu et al). 1996;17:7-13
Species-specific in situ hybridization with fluorochrome-labeled DNA probes to study vascularization of human skin grafts on athymic mice (Young et al). 1996;17:305-10
The use of skin from a monozygotic twin combined with cultured epithelial autografts as coverage for a large surface area burn: a case report and review of the literature (Caruso et al). 1996;17:432-4 (Case rep.)
Transplantation, homologous
Clinical evaluation of an acellular allograft dermal matrix in full-thickness burns (Wainwright et al). 1996;17:124-36
Growth peptide release from biologic dressings: a comparison (Rennekampff et al). 1996;17:522-7

Transplantation, homologous—cont'd
Integrin and matrix molecule expression in cultured skin replacements (Rennekampff et al). 1996;17:213-21
A reliable and cost-effective in vitro assay of skin viability for skin banks and burn centers (Klein et al). 1996; 17:565-70
Trifluoperazine
Calcium antagonists alter cell shape and induce procollagenase synthesis in keloid and normal human dermal fibroblasts (Doong et al). 1996;17:497-514

U

Ulnar nerve
Elbow anchylosis due to postburn heterotopic ossification (Holguin et al). 1996;17:150-4
Unna boot dressing
Unna's boot dressings facilitate outpatient skin grafting of hands (Sanford and Gore). 1996;17:323-6

V

Verapamil
Calcium antagonists alter cell shape and induce procollagenase synthesis in keloid and normal human dermal fibroblasts (Doong et al). 1996;17:497-514
Vimentin
Progressive burn injury documented with vimentin immunostaining (Nanney et al). 1996;17:191-8
Vital statistics
Burn incidence and medical care use in the United States: estimates, trends, and data sources (Brigham and McLoughlin). 1996;17:95-107
Vitronectin
The effect of an arginine–glycine–aspartic acid peptide and hyaluronate synthetic matrix on epithelialization of meshed skin graft interstices (Cooper et al). 1996; 17:108-16

W

Washing; *see* **Baths**
World Health Organization
Burns of the world (Munster). 1996;17:477-84
Wound healing
Assessment of cosmetic and functional results of conservative versus surgical management of facial burns (Fraulin et al). 1996;17:19-29
Effects of an arginine–glycine–aspartic acid peptide-containing artificial matrix on epithelial migration in vitro and experimental second-degree burn wound healing in vivo (Mertz et al). 1996;17:199-206
Essential microminerals and their response to burn injury (Gamliel et al). 1996;17:264-72
Growth peptide release from biologic dressings: a comparison (Rennekampff et al). 1996;17:522-7
Hyaluronic acid impedes reepithelialization of skin graft donor sites (Bettinger et al). 1996;17:302-4
Pulsed electromagnetic fields in experimental cutaneous wound healing in rats (Patiño et al). 1996;17:528-31
Substance P has a role in neurogenic mediation of human burn wound healing (Dunnick et al). 1996;17: 390-6
Wound coverage: Is there a difference? (Hastings et al). 1996;17:416-20
Wounds and injuries
Propane gas dangers and strategies for prevention of injuries (Smoot and Hickerson). 1996;17:273-9
Wrist
Unna's boot dressings facilitate outpatient skin grafting of hands (Sanford and Gore). 1996;17:323-6